

Properties of the $\Lambda(1405)$ Measured at CLAS

Kei Moriya and Reinhard Schumacher
for the CLAS collaboration

Department of Physics, Carnegie Mellon University, Pittsburgh, PA 15213, USA

Abstract

Using the CLAS detector system in Hall B at Jefferson Lab, we have done a high statistics measurement of the photoproduction of $K^+ \Lambda(1405)$ using a proton target. The reconstructed invariant mass seen in the various $\Sigma\pi$ decay modes has been measured, as well as the differential cross section $\frac{d\sigma}{d\tau}$. The nature of the $\Lambda(1405)$ with its peculiar mass distribution is not well understood, and studies have shown that it may possess strong dynamical components, which could be extracted from our data. Various aspects of our analysis are discussed, as well as future prospects to further enhance the precision of the results.

Key words: hyperons, photoproduction, CLAS, $\Lambda(1405)$

PACS: 14.20.Jn, 25.20.Lj

1. Introduction

The $\Lambda(1405)$ is a well-established hyperon resonance located just below $N\bar{K}$ threshold. Even though its existence has been known since the 1960's and it holds a four-star rating in the PDG [1], its nature as a resonance is not well understood. The $\Lambda(1405)$ can only be seen by reconstructing the invariant mass of the $\Sigma\pi$ system into which it decays. Previous experiments [2, 3] have shown this mass distribution, or “lineshape” to be distorted from the usual Breit-Wigner resonance form, leading to various speculations on the nature of the $\Lambda(1405)$.

Using the CLAS [4] detector system in Hall B at the Thomas Jefferson National Accelerator Laboratory, we have done a photoproduction measurement of the $\Lambda(1405)$. Due to the high statistics of the experiment and good resolution of the CLAS system, we were able to extract the lineshapes and differential cross sections of the $\Lambda(1405)$ in all three of its $\Sigma\pi$ decay modes. The lineshapes and cross sections will give a much improved quantitative description of the $\Lambda(1405)$, and will lead to further insights into its nature and production mechanism.

2. Theory of the $\Lambda(1405)$ Mass Distribution

In recent years, theoretical progress in chiral dynamics in the $S = -1$ sector has gained much attention, and has been extended to include a description of the $\Lambda(1405)$. In this approach the $\Lambda(1405)$ is dynamically generated by iterating over the meson-baryon interactions among the ground state baryon octet and pseudoscalar mesons octet. A paper by Nacher *et al.* [5] predicts, using a chiral unitary approach, the lineshapes of the $\Lambda(1405)$ as it decays to its three $\Sigma\pi$ decay modes. Figure 1 shows the predicted lineshapes, a result of the isospin 0 and 1 amplitudes




Figure 1: Theoretical lineshapes predicted [5] for the $\Lambda(1405)$ using the chiral unitary approach. Due to the interfering isospin 0 and 1 terms, the lineshapes are distorted from a simple Breit-Wigner form. In more recent years, attention has also focused on the possible two pole structure of the $\Lambda(1405)$ in the $I = 0$ channel alone. This too is expected to influence the lineshapes [6].

interfering to produce shifts in the $\Sigma^+\pi^-$ and $\Sigma^-\pi^+$ lineshapes in opposite directions. Unlike an ordinary resonance where the lineshape will follow a simple Breit-Wigner form, the $\Lambda(1405)$ spectrum is predicted to be strongly distorted due to a mixture of contributions. This figure has been the starting point in our analysis, and our work has been focused on comparing our experimental data with this prediction.

3. Data Analysis Using CLAS

In this section we show some of the work that was done at CLAS to study the photoproduction of the $\Lambda(1405)$. The reaction of interest at CLAS is $\gamma + p \rightarrow K^+ + \Lambda(1405)$, with the $\Lambda(1405)$ decaying exclusively into $\Sigma\pi$. The CLAS data set was obtained in 2004 using tagged real photons up to 3.84 GeV, and a two charged track trigger. For the $\Sigma^+\pi^-$ decay mode, there are two possible decays for the Σ^+ , i.e. to $p\pi^0$ or $n\pi^+$. The first step of the analysis is to select either (p, K^+, π^-) or (K^+, π^+, π^-) events.

After the three charged particles are selected and identified using momentum and TOF information, the missing mass of the event is calculated to obtain the missing particle(s). Figure 2(a) shows an example of the missing mass squared (MM^2) for one bin of our analysis. A Gaussian fit to the neutron peak had a $\pm 3\sigma$ cut applied. Next, we combine our final state particles to reconstruct the intermediate ground state hyperon that the reaction went through. In Figure 2(b) we show the result of plotting the invariant masses squared of the (n, π^+) versus (n, π^-) . Again a fit was done and a region of $\pm 2\sigma$ was selected around each peak; the Σ overlap region was excluded as well.

At this stage we are ready to fit our resulting $\Sigma\pi$ spectrum as a sum of templates of Monte Carlo (MC) spectra. The MC were initially generated with mass and width values given in the PDG [1] and processed through the standard CLAS simulation package to account for detection inefficiencies. After this fitting process, the non- $\Lambda(1405)$ contributions were subtracted off from the data and the “residual” spectrum was taken to be from the $\Lambda(1405)$. The resulting lineshape was then acceptance corrected bin by bin in mass. The final lineshape spectrum was then obtained by summing over all the production angle bins in each energy bin. Our final lineshapes are thus shown this way, acceptance corrected and summed over all production angles.

Once the above process of fitting and obtaining the lineshapes in each energy bin was completed, the resulting lineshapes were then used as input to generate MC templates for the $\Lambda(1405)$, so that the process could be iterated and refined. The fit result in one sample bin is shown in Figure 3(a) for the iterated version.




Figure 2: (Color online) (a) The missing mass squared (MM^2) calculated for one energy and angle bin for the reaction $\gamma + p \rightarrow K^+ + \pi^+ + \pi^- + X$, where X denotes the missing particle(s). The neutron peak is clearly seen. The fit result of a Gaussian (red line) and a polynomial background (green line) are also shown. The $\pm 3\sigma$ range of the fit is shown with the red dashed lines. (b) $M^2(n, \pi^+)$ vs $M^2(n, \pi^-)$ for a particular bin when detecting K^+, π^+, π^- and reconstructing the missing neutron. The horizontal band corresponds to Σ^+ events, and the vertical band to Σ^- events. A faint diagonal band of neutral kaons decaying to π^+, π^- is also seen. $\pm 2\sigma$ selection cuts are shown for each hyperon.

4. Results

After the iteration of the fit, the mass distribution we obtained is shown in Figure 3(b) for one energy bin near threshold of the $\Lambda(1405)$. Our lineshape result is shown for the three different charge combinations $\Sigma^+\pi^-$ (red), $\Sigma^0\pi^0$ (blue), and $\Sigma^-\pi^+$ (green). The $\Sigma^+\pi^-$ spectrum is a weighted average of the two different decay channels of the Σ^+ . The difference in the lineshapes is clear, with the $\Sigma^+\pi^-$ spectrum peaking at the highest mass, and having a much sharper structure than the $\Sigma^-\pi^+$ spectrum.

A comparison of our experimental result can be done with the theoretical prediction (Figure 1). Although the lineshapes differ from each other as predicted, we see that for the theory prediction, the $\Sigma^-\pi^+$ spectrum is shifted toward the higher mass, whereas this applies to the $\Sigma^+\pi^-$ spectrum in our experimental result. The difference in the two charged decay modes, according to the theory, is based on an interference between isospin $I = 0$ and $I = 1$ terms, as shown in [5]. At the moment, this “interchange” of lineshapes is an unresolved issue that we hope to resolve in future studies.

5. Future Prospects and Conclusion

The mass distributions of the $\Lambda(1405)$ have been obtained in nine different energy bins, and although much work has been put into obtaining these, there is still room for improvement. One example of prospective work is using the kinematic fitter that was developed for CLAS. The results shown here do not utilize the kinematic fitter, but our preliminary studies show that our resolution is readily improved when using it.

Preliminary differential cross sections for the $\Lambda(1405)$ were also shown during the presentation, which were obtained by summing over each lineshape in each energy and angle bin. The values of this $\frac{d\sigma}{dt}$ showed differences between the different charge combinations, not just in scale but also in dependence on t . The nature of this different t -dependence for each decay mode is unknown.




Figure 3: (Color online) (a) Sample of fit to $M(\Sigma\pi)$ spectrum in one bin of the $\Sigma^+ \rightarrow n\pi^+$ decay mode. Only the heights of the MC templates were allowed to change. The $\Sigma(1385)$ yield was measured in the $\Lambda\pi^0$ decay mode, and scaled by branching ratio into the $\Sigma\pi$ decay modes. This figure is for the iterated version of the fit, see text for details. (b) The final $\Lambda(1405)$ lineshape result that we obtained for one energy bin near threshold. The three different lineshapes are plotted against each other. Stark differences are observed, especially between the two charged decay modes. The dashed line is a relativistic Breit-Wigner shape using the nominal values of mass and width of the $\Lambda(1405)$ given in the PDG. Notice that although the $\Sigma^+\pi^-$ threshold is lower than that of $\Sigma^-\pi^+$, the $\Sigma^-\pi^+$ spectrum appears at a lower mass.

Another prospect for our analysis is direct experimental evidence for the spin and parity of the $\Lambda(1405)$. Although the majority of theories assume $J^P = \frac{1}{2}^-$, no direct experimental proof for these quantum numbers has been given [7]. Our preliminary analysis shows a detectable polarization for the $\Lambda(1405)$, which is transferred to the Σ^+ , allowing us to infer its spin and parity. Our results are consistent with the common expectation, but will nonetheless be an important step in identifying the nature of the $\Lambda(1405)$. Also, it should be noted that from these measurements the polarization of the $\Lambda(1405)$ is measured, giving further information about its production mechanism.

In conclusion, an overview of the CLAS analysis of the $\Lambda(1405)$ has been given. We have an unprecedented high statistics data set for the $\Lambda(1405)$, and have obtained the lineshapes and differential cross sections in energy bins spanning from threshold up to 2.84 GeV in the overall center of mass energy. We observe a difference in the different decay mode lineshapes, leading to a difference in the differential cross sections. We have just started comparing our results to theory, and are eager to finalize our results in the coming months.

References

- [1] C. Amsler *et al.* [Particle Data Group], Phys. Lett. B **667**, 1 (2008).
- [2] D. W. Thomas, A. Engler, H. E. Fisk and R. W. Kraemer, Nucl. Phys. B **56**, 15 (1973).
- [3] R. J. Hemingway, Nucl. Phys. B **253**, 742 (1985).
- [4] B. A. Mecking *et al.* [CLAS Collaboration], Nucl. Instrum. Meth. A **503**, 513 (2003).
- [5] J. C. Nacher, E. Oset, H. Toki and A. Ramos, Phys. Lett. B **455**, 55 (1999) [arXiv:nucl-th/9812055].
- [6] D. Jido, J. A. Oller, E. Oset, A. Ramos and U. G. Meissner, Nucl. Phys. A **725**, 181 (2003) [arXiv:nucl-th/0303062].
- [7] An analysis of the spin of the $\Lambda(1405)$ was given in A. Engler, H. E. Fisk, R. W. Kraemer, C. M. Meltzer and J. B. Westgard, Phys. Rev. Lett. **15**, 224 (1965), which concluded that “spin greater than $\frac{1}{2}$ is not required by our data.” $J^P = \frac{1}{2}^+$ is considered in O. Kittel *et al.*, arXiv:hep-ph/0010186, arXiv:hep-ph/0508150.